

## LITERATUR

### 11.1 TERMINOLOGIE

Dilling H, Mombour W, Schmidt MH (1991). Internationale Klassifikation psychischer Störungen – ICD-10. Kapitel V (F): Klinisch-diagnostische Leitlinien. WHO 1. A. Bern, Göttingen, Toronto: Huber.

### 11.2 EPIDEMIOLOGIE UND VERLAUF

- Alonso J, Angermeyer MC, Bernert S, et al. (2004) for the ESEMeD/MHEDEA 2000 Investigators, European Study of the Epidemiology of Mental Disorders (ESEMeD) Project. Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatr Scand* (Suppl 420): 21–27.
- Angst J (1987). Begriff der affektiven Erkrankungen. In: Urscher KP, Lauber H, Meyer J-E et al. (Hrsg.). *Psychiatrie der Gegenwart*. Bd. 5: Affektive Psychosen. Berlin, Heidelberg, New York: Springer, S. 51–68.
- Bostwick JM, Pankratz VS (2000). Affective disorders and suicide risk: a reexamination. *Am J Psychiatry* 157: 1925–1932.
- Dilling H (Hrsg.) (2002). Lexikon zur ICD-10-Klassifikation psychischer Störungen. Bern, Göttingen, Toronto: Huber.
- Dunner DL, Schmaling KB, Hendrickson H, et al. (1996). Cognitive therapy versus fluoxetine in the treatment of dysthymic disorder. *Depression* 4: 34–41.
- Eaton WW, Shao H, Nestadt G, et al. (2008). Population-based study of first onset and chronicity in major depressive disorder. *Arch Gen Psychiatry* 65: 513–520.
- Goodwin FK, Jamison KR (2007). *Manic-Depressive Illness*. 2<sup>nd</sup> ed. Oxford, New York, Tokyo: Oxford University Press.
- Jacobi F, Wittchen H-U, Höltig C, et al. (2004). Prevalence, co-morbidity and correlates of mental disorders in the general population: results from the German Health Interview and Examination Survey (GHS). *Psychol Med* 34: 579–611.
- Keller MB, McCullough JP, Klein DN, et al. (2000). A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *N Engl J Med* 342: 1462–1470.
- McCullough JP (2003). Treatment for chronic depression using Cognitive Behavioral Analysis System of Psychotherapy (CBASP). *J Clin Psychol* 59: 833–846.
- Murray CJL, Lopez AD (1996). The Global Burden of Disease: A comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Geneva: World Health Organization.
- Thase ME, Rush AJ (1995). Treatment resistant depression. In: Bloom FE, Kupfer DJ (eds.). *Psychopharmacology: The Fourth Generation of Progress*. New York: Raven Press, pp. 1081–1097.
- Vos T, Barber RM Bell B, et al. (2015). Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 386(9995): 743–800.

### 11.3 SYMPTOMATIK UND TYPISIERUNG

- Angst J, Merikangas K, Scheidegger P, Wicki W (1990). Recurrent brief depression: a new subtype of affective disorder. *J Affect Disord* 19: 87–98.
- APA – American Psychiatric Association (1994). *Diagnostic and Statistical Manual of Mental Disorders – DSM-IV*. Washington, DC: American Psychiatric Association.
- APA – American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders – DSM-5*. Washington, DC: American Psychiatric Association.
- APA – American Psychiatric Association (2015). *Diagnostisches und Statistisches Manual Psychischer Störungen DSM-5*. Deutsche Ausgabe hrsg. von P. Falkai und H-U Wittchen. Göttingen, Bern, Wien: Hogrefe.
- Bostwick JM, Pankratz VS (2000). Affective disorders and suicide risk: a reexamination. *Am J Psychiatry* 157(12): 1925–1932.

- Dunner DL (2001). Acute and maintenance treatment of chronic depression. *J Clin Psych* 62: 10–16.
- Klerman GL, Endicott R, Spiker RM, Hirschfeld A (1979). Neurotic depression: a systematic analysis of multiple criteria of meanings. *Am J Psychiatry* 136: 57–61.
- Łojko D, Rybakowski JK (2017). Atypical depression: current perspectives. *Neuropsychiatr Dis Treat* 13: 2447–2456.
- McCullough JP (2000). *Treatment for Chronic Depression: Cognitive Behavioral Analysis System of Psychotherapy (CBASP)*. New York: Guilford Press.
- Moussavi S, Chatterji S, Verdes E, et al. (2007). Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet* 370: 851–858.
- Westrin A, Lam RW (2007). Seasonal affective disorder: a clinical update. *Ann Clin Psychiatry* 19: 239–246.
- WHO – Weltgesundheitsorganisation (1991). Internationale Klassifikation psychischer Störungen, ICD-10, Kapitel V (F). Bern, Göttingen, Toronto: Huber.
- Winokur G, Clayton P, Reich T (1969). *Manic Depressive Illness*. St. Louis, MO: Mosby.

### 11.4 ÄTILOGIE UND PATHOGENESE

- Bibring E (1953). Das Problem der Depression. *Psyche* 6: 81–101.
- Beck AT (1976). *Cognitive Therapy and the Emotional Disorders*. New York: International Universities Press.
- Berger M, van Calker D, Riemann D (2003). Sleep and manipulations of the sleep–wake rhythm in depression. *Acta Psychiatr Scand* 108: 83–91.
- Bertelsen A, Harvald B, Hauge M (1977). A Danish twin study of manic-depressive disorders. *Br J Psychiatry* 130: 330–351.
- Binneman B, Feltner D, Kolluri S, et al. (2008). A 6-week randomized, placebo-controlled trial of CP-316,311 (a selective CRH1 antagonist) in the treatment of major depression. *Am J Psychiatry* 165(5): 617–620.
- Bohus M, Berger M (1992). Der Beitrag biologisch-psychiatrischer Befunde zum Verständnis depressiver Erkrankungen. *Z Klin Psychol* 21: 156–171.
- Bowlby J (1988). Developmental psychiatry comes to age. *Am J Psychiatry* 145: 1–10.
- Brakemeier EL, Hautzinger M (2008). *Kognitive Verhaltenstherapie (KVT)*. In: Bschor T (Hrsg.). *Behandlungsmanual therapieresistente Depression*. Pharmakotherapie – somatische Therapieverfahren – Psychotherapie. Stuttgart: Kohlhammer, S. 330–367.
- Brakemeier EL, Dobias J, Hertel J, et al. (2018). Childhood maltreatment in women with borderline personality disorder, chronic depression, episodic depression, and in healthy controls. *Psychother Psychosom* 87: 49–51.
- Bunney WE Jr, Davis JM (1965). Norepinephrine in depressive reactions. A review. *Arch Gen Psychiatry* 13: 483–494.
- Dozier M, Stovall-McCough KC, Albus KE (1999). Attachment and psychopathology in adulthood. In: Cassidy J, Shaver P (eds.). *Handbook of Attachment: Theory, research, and clinical applications*. New York: Guilford Press, pp. 497–519.
- Dreßing H, Riemann D (1994). *Diagnostik und Therapie von Schlafstörungen*. Stuttgart, Jena, New York: Fischer.
- Freud S (1917). *Trauer und Melancholie*. GW Bd. X. Frankfurt.
- Furey ML, Drevets WC (2006). Antidepressant efficacy of the antimuscarinic drug scopolamine: a randomized, placebo-controlled clinical trial. *Arch Gen Psychiatry* 63(10): 1121–1129.
- Gilbert P (1984). *Depression. From Psychology to Brain State*. London, Hillsdale, NJ: Lawrence Erlbaum.
- Goodwin FK, Jamison KR (2007). *Manic-Depressive Illness*. 2<sup>nd</sup> ed. Oxford, New York, Tokyo: Oxford University Press.
- Hoffmann SO, Hochapel G (2009). *Neurosenlehre, psychotherapeutische und psychosomatische Medizin*. 8. überarb. und erw. A. Stuttgart, New York: Schattauer.
- Janowsky DS, el Yousef MK, Davis JM, Sekerke HJA (1972). A cholinergic-adrenergic hypothesis of mania and depression. *Lancet* 23: 632–635.
- Lewinsohn PM, Youngren MA, Grosscup SJ (1979). Reinforcement and depression. In: Depue RA (ed.). *The Psychobiology of Depressive Disorders*. New York: Academic Press, pp. 291–319.

## 444.e2 11 Affektive Störungen

- Li N, Lee B, Liu RJ, et al. (2010). mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science* 329(5994): 959–964.
- Manji HK, Duman RS (2001). Impairments of neuroplasticity and cellular resilience in severe mood disorders: implications for the development of novel therapeutics. *Psychopharmacol Bull* 35: 5–49.
- Mill J, Petronis A (2007). Molecular studies of major depressive disorder: the epigenetic perspective. *Mol Psychiatry* 12(9): 799–814.
- Modell S, Ising M, Holsboer F, Lauer CJ (2005). The Munich vulnerability study on affective disorders: premorbid polysomnographic profile of affected high-risk probands. *Biol Psychiatry* 58(9): 694–699.
- Pietrek C, Elbert T, Weierstall R, et al. (2013). Childhood adversities in relation to psychiatric disorders. *Psychiatry Res* 206: 103–110.
- Riemann D, Hohagen F, Bahro M, Berger M (1994). Sleep in depression: the influence of age, gender and diagnostic subtype on baseline sleep and the cholinergic REM induction test with RS86. *Eur Arch Psychiatry Clin Neurosci* 243: 279–290.
- Royal K, Theobold D, Graham A, et al. (2007). Mania-like behavior induced by disruption of CLOCK. *Proc Natl Acad Sci USA* 104(15): 6406–6411.
- Schramm E (2011). Interpersonelle Psychotherapie bei Depressionen und anderen psychischen Störungen. Stuttgart, New York: Schattauer.
- Seligman MEP (1975). Helplessness. San Francisco: Freeman.
- Serchov T, Clement HW, Schwarz MK, et al. (2015). Increased signaling via adenosine a<sub>1</sub> receptors, sleep deprivation, imipramine, and ketamine inhibit depressive-like behavior via induction of Homer1a. *Neuron* 87(3): 549–562.
- Serchov T, Heumann R, van Calker D, Biber K (2016). Signaling pathways regulating Homer1a expression: implications for antidepressant therapy. *Biol Chem* 397(3): 207–214.
- Schloesser RJ, Martinowich K, Manji HK (2012). Mood-stabilizing drugs: mechanisms of action. *Trends Neurosci* 35(1): 36–46.
- Stankov L (2013). Depression and life satisfaction among European and Confucian adolescents. *Psychol Assess* 25: 1220–1234.
- Stenzel N, Bollmann S, Brakemeier EL (2017). Depression. In: Brakemeier EL, Jacobi F (Hrsg.). Verhaltenstherapie in der Praxis. Weinheim: Beltz, S. 500–512.
- Tellenbach H (1961). Melancholie. Berlin, Heidelberg, New York: Springer.
- Tsankova N, Renthal W, Kumar A, Nestler EJ (2007). Epigenetic regulation in psychiatric disorders. *Nat Rev Neurosci* 5: 355–367.
- van Calker D, Serchov T, Normann C, Biber K (2018). Recent insights into antidepressant therapy: distinct pathways and potential common mechanisms in the treatment of depressive syndromes. *Neurosci Biobehav Rev* 88: 63–72.
- von Zerssen D, Berger M, Dose M, et al. (1986). The nature of neuroendocrine abnormalities in depression: a controversial issue in contemporary psychiatry. *Psychiatr Develop* 3: 237–256.
- Wiersam JE, Hovens JGFM, van Oppen P, et al. (2009). The importance of early trauma and childhood life events for chronicity of depression. *J Clin Psychiatry* 70: 983–989.
- Wu JC, Bunney WE (1990). The biological basis of an antidepressant response to sleep deprivation and relapse: a review and hypothesis. *Am J Psychiatry* 147: 14–21.
- Zarate C Jr, Machado-Vieira R, Henter I, et al. (2010). Glutamatergic modulators: the future of treating mood disorders? *Harv Rev Psychiatry* 5: 293–303.

## 11.5 DIFFERENZIALDIAGNOSTISCHER PROZESS

- APA – American Psychiatric Association (1994). Diagnostic and Statistical Manual of Mental Disorders – DSM-IV. Washington, DC: American Psychiatric Association.
- Soreft SM, McNeil GN (1987). Handbook of Psychiatric Differential Diagnosis. Littleton: PSG Publishing Company.
- Zong S, Hoffmann C, Mané-Damas M, et al. (2017). Neuronal surface auto-antibodies in neuropsychiatric disorders: Are there implications for depression? *Front Immunol* 8: 752.

## 11.6 AKUTTHERAPIE AFFEKTIVER ERKRANKUNGEN

- Adli M, Bauer M, Rush AJ (2006). Algorithms and collaborative-care systems for depression: are they effective and why? A systematic review. *Biol Psychiatry* 59(11): 1029–1038.
- Allen MH, Hirschfeld RM, Wozniak PJ, et al. (2006). Linear relationship of valproate serum concentration to response and optimal serum levels for acute mania. *Am J Psychiatry* 163(2): 272–275.
- Altshuler LL, Bauer M, Frye MA (2001). Does thyroid supplementation accelerate tricyclic antidepressants response: a review and meta-analysis. *Am J Psychiatry* 158: 1617–1622.
- Amick HR, Gartlehner G, Gaynes BN, et al. (2015). Comparative benefits and harms of second generation antidepressives and cognitive behavioral therapies in initial treatment of major depressive disorder: systematic review and metaanalysis. *BMJ* 351: h6019.
- Anderson IM (2000). Selective serotonin reuptake inhibitors versus tricyclic antidepressants: a meta-analysis of efficacy and tolerability. *J Affect Disord* 58(1): 19–36.
- Andrade C, Sandarsh S, Chethan KB, Nagesh KS (2010). Serotonin reuptake inhibitor antidepressants and abnormal bleeding: a review for clinicians and a reconsideration of mechanisms. *J Clin Psychiatry* 71(12): 1565–1575.
- APA – American Psychiatric Association (2001). The Practice of Electroconvulsive Therapy. Recommendations for Treatment, Training and Privileging: A Task Force Report of the American Psychiatric Association. 2<sup>nd</sup> ed. Washington, DC.
- Arnow BA, Constantino MJ (2003). Effectiveness of psychotherapy and combination treatment for chronic depression. *J Clin Psychol* 59: 893–905.
- Baldessarini RJ, Tondo L, Hennen J (2003). Lithium treatment and suicide risk in major affective disorders. *J Clin Psychiatry* 63 (Suppl 5): 44–52.
- Barth J, Munder T, Gerger H, et al. (2013). Comparative efficacy of seven psychotherapeutic interventions for patients with depression: a network meta-analysis. *PLoS Med* 10(5): e1001454.
- Bauer M, Dopfner S (1999). Lithium augmentation in treatment-resistant depression: meta-analysis of placebo-controlled studies. *J Clin Psychopharmacol* 19: 427–434.
- Beck AT, Rush AJ, Shaw BF, Emery G (1979). Cognitive Therapy of Depression. New York: Guilford Press.
- Bell AC, D'Zurilla TJ (2009). Problem-solving therapy for depression: a meta-analysis. *Clin Psychol Rev* 29(4): 348–353.
- Benedetti G (1987). Analytische Therapie der affektiven Psychosen. In: Kisker KP, Lauter H, Meyer J-E et al. (Hrsg.). Psychiatrie der Gegenwart. Bd. 5: Affektive Psychosen. 3. A. Berlin, Heidelberg, New York: Springer, S. 369–386.
- Benedetti F, Barbini B, Campori E, et al. (2001). Sleep phase and lithium to sustain the antidepressant effect of total sleep deprivation in bipolar depression: new findings supporting the internal coincidence model? *J Psychiatr Res* 35: 323–329.
- Berger M, Riemann D, Wiegand M, et al. (1993). Polysomnographische Beiträge zum Verständnis der Pathogenese affektiver Erkrankungen. In: Berger M, Möller H-J, Wittchen H-U (Hrsg.). Psychiatrie als empirische Wissenschaft. München, Bern, Wien, New York: Zuckschwerdt, S. 128–140.
- Beutel ME, Leuzinger-Bohleber M, Rüger B, et al. (2012). Psychoanalytic and cognitive-behavior therapy of chronic depression: study protocol for a randomized controlled trial. *Trials* 13: 117.
- Brakemeier EL, Merkl A, Wilbertz G, et al. (2014). Cognitive-behavioral therapy as continuation treatment to sustain response after electroconvulsive therapy in depression: a randomized controlled trial. *Biol Psychiatry* 76(3): 194–202.
- Brakemeier EL, Radtke M, Engel V, et al. (2015). Overcoming treatment resistance in chronic depression: a pilot study on outcome and feasibility of the cognitive behavioral analysis system of psychotherapy as an inpatient treatment program. *Psychother Psychosom* 84(1): 51–56.
- Brakemeier EL, Zimmermann J, Erz E et al. (2017). Interpersonelles Integratives Modellprojekt für Geflüchtete mit psychischen Störungen. *Psychotherapeut* 62(4): 322–332.

- Brakemeier EL, Dobias J, Hertel J, et al. (2018). Childhood maltreatment in women with borderline personality disorder, chronic depression, episodic depression, and in healthy controls. *Psychother Psychosom* 87: 49–51.
- Browne G, Steiner M, Roberts J, et al. (2002). Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs. *J Affect Disord* 68: 317.
- Bschor T (2008). Behandlungsmanual Therapieresistente Depression. Stuttgart: Kohlhammer.
- Calabrese JR, Keck PE Jr, Macfadden W, et al. (2005a). A randomized, double-blind, placebo-controlled trial of quetiapine in the treatment of bipolar I or II depression. *Am J Psychiatry* 162(7): 1351–1360.
- Calabrese JR, Shelton MD, Rapoport DJ, et al. (2005b). A 20-month, double-blind, maintenance trial of lithium versus divalproex in rapid-cycling bipolar disorder. *Am J Psychiatry* 162(11): 2152–2161.
- Calabrese JR, Huffman RF, White RL, et al. (2008). Lamotrigine in the acute treatment of bipolar depression: results of five double-blind, placebo controlled clinical trials. *Bipolar Disord* 10: 323–333.
- Casper F (2017). Evidenzbasierte Psychotherapie im klinischen Alltag. *Info Neurologie & Psychiatrie* 19(1): 43–50.
- Cipriani A, Hawton K, Stockton S, Geddes JR (2013). Lithium in the prevention of suicide in mood disorders: updated systematic review and meta-analysis. *BMJ* 346: f3646.
- Clarke GN, Rohde P, Lewinsohn PM, et al. (1999). Cognitive-behavioral treatment of adolescent depression: efficacy of acute group treatment and booster sessions. *J American Academy of Child & Adolescent Psychiatry*, 38(3), 272–279.
- Cordoba S, de Asis J, Kotbi N, Mahgoub N (2011). Worsening mania associated with slow increase of quetiapine dose. *J Neuropsychiatry Clin Neurosci* 23: E5–6.
- Crits-Christoph P (1992). The efficacy of brief dynamic psychotherapy: a meta-analysis. *Am J Psychiatry* 149(2): 151–158.
- Cuijpers P (2016). Are all psychotherapies equally effective in the treatment of adult depression? The lack of statistical power of comparative outcome studies. *Evid Based Ment Health* 19(2): 39–42.
- Cuijpers P (2017). Four decades of outcome research on psychotherapies for adult depression: an overview of a series of metaanalyses. *Can Psychol* 58: 7–19.
- Cuijpers P, van Straten A, Warmerdam L (2007). Problem solving therapies for depression: a meta-analysis. *Eur Psychiatry* 22: 9–15.
- Cuijpers P, van Straten A, van Oppen P, Andersson G (2008). Are psychological and pharmacologic interventions equally effective in the treatment of adult depressive disorders? A meta-analysis of comparative studies. *J Clin Psychiatry* 69(11): 1675–1685.
- Cuijpers P, van Straten A, Warmerdam L, Andersson G (2009). Psychotherapy versus the combination of psychotherapy and pharmacotherapy in the treatment of depression: a meta-analysis. *Depress Anxiety* 26(3): 279–288.
- Cuijpers P, van Straten A, Schuurmans J, et al. (2010). Psychotherapy for chronic major depression and dysthymia: a meta-analysis. *Clin Psychol Rev* 30(1): 51.
- Cuijpers P, Hollon SD, van Straten A, et al. (2013). Does cognitive behaviour therapy have an enduring effect that is superior to keeping patients on continuation pharmacotherapy? A meta-analysis. *BMJ Open* 3(4). pii: e002542.
- Cuijpers P, Donker T, Weissman MM, et al. (2016). Interpersonal Psychotherapy for mental health problems: a comprehensive meta-analysis. *Am J Psychiatr* 173: 680–687.
- David D, Cristea I, Hofmann SG (2018). Why cognitive behavioral therapy is the current gold standard of psychotherapy. *Front Psychiatry* 9: 4. doi: 10.3389/fpsy.2018.00004. eCollection 2018doi:10.3389/fpsy.2018.0004.
- Davis MP, Khawam E, Pozuelo L, et al. (2002). Management of symptoms associated with advanced cancer: olanzapine and mirtazapine. A World Health Organization project. *Expert Rev Anticancer Ther* 2: 365–376.
- DeBattista C, Solvason B, Poirier J, et al. (2005). A placebo-controlled, randomized, double-blind study of adjunctive bupropion sustained release in the treatment of SSRI-induced sexual dysfunction. *J Clin Psychiatry* 66(7): 844–848.
- de Bodinat C, Guardiola-Lemaitre B, Mocaer E, et al. (2010). Agomelatine, the first melatonergic antidepressant: discovery, characterization and development. *Nat Rev Drug Discov* 9(8): 628–642.
- de Maat S, de Jonghe F, de Kraker R, et al. (2013). The current state of the empirical evidence for psychoanalysis: a meta-analytic approach. *Harvard Rev Psychiatry* 21(3): 107–137.
- DeRubeis RJ, Gelfand LA, Tang TZ, et al. (1999). Medications versus cognitive behavior therapy for severely depressed outpatients: mega-analysis of four randomized comparisons. *Am J Psychiatry* 156(7): 1007–1013.
- Driessens E, Cuijpers P, de Maat S, et al. (2010). The efficacy of short-term psychodynamic psychotherapy for depression: a meta-analysis. *Clin Psychol Rev* 30(1): 25–36.
- Driessens E, Hegelmaier LM, Abbass AA, et al. (2015). The efficacy of short-term psychodynamic psychotherapy for depression: a meta-analysis update. *Clin Psychol Rev* 42: 1–15.
- Dubovsky S, Buzan R (1999). Mood disorders. In: Hales R, Yudofsky SC, Talbott JA (eds.). *Textbook of Psychiatry*. Washington: American Psychiatric Press, pp. 479–567.
- D'Zurilla TJ, Goldfried MR (1971). Problem solving and behavior modification. *J Abnormal Psychol* 78: 78(1): 107–126.
- Edwards JG, Anderson I (1999). Systematic review and guide to selection of selective serotonin reuptake inhibitors. *Drugs* 57: 507–533.
- Elliott R, Greenberg LS, Lietaer G (2004). Research on experiential psychotherapies. In: Lambert MJ (ed.). *Bergin and Garfield's Handbook of Psychotherapy and Behavior Change*. 5<sup>th</sup> ed. New York: Wiley, pp. 493–540.
- Eyding D, Lelgemann M, Grouven U, et al. (2010). Problem solving and behaviour. Reboxetine for acute treatment of major depression: systematic review and meta-analysis of published and unpublished placebo and selective serotonin reuptake inhibitor controlled trials. *BMJ* 341: c4737.
- Geddes J, Butler R (2002). Depressive disorders. In: BMJ Publishing Group (ed.). *Clinical Evidence. Mental Health. The International Source of the Best Available Evidence for Mental Health Care*. Margate, Kent: Thanet Press, p. 6176.
- Gellis ZD, McGinty J, Horowitz A, et al. (2007). Problem-solving therapy for late-life depression in home care: a randomized field trial. *Am J Geriatr Psychiatry* 11: 968–978.
- George MS, Taylor JJ, Short B (2013). Treating the depressions with superficial brain stimulation methods. *Handb Clin Neurol* 116: 399–413.
- Gerber AJ, Kocsis JH, Milrod BL, et al. (2011). A quality-based review of randomized controlled trials of psychodynamic psychotherapy. *Am J Psychiatry* 168: 19–28.
- Geschwind N, Peeters F, Huibers M, et al. (2012). Efficacy of mindfulness-based cognitive therapy in relation to prior history of depression: randomised controlled trial. *Br J Psychiatry* 201(4): 320–325.
- Gloaguen V, Cottraux J, Cucherat M, Blackburn IM (1998). A meta-analysis of the effects of cognitive therapy in depressed patients. *J Affect Disord* 49: 59–72.
- Greenberg LS, Rice LN, Elliott R (1993). *Facilitating Emotional Change. The moment-by-moment process*. New York: Guilford Press.
- Guidi J, Tomba E, Fava GA (2016). The sequential integration of pharmacotherapy and psychotherapy in the treatment of major depressive disorder: a meta-analysis of the sequential model and a critical review of the literature. *Am J Psychiatry* 173: 128–137.
- Hallahan B, Hibbeln JR, Davis JM, et al. (2007). Omega-3 fatty acid supplementation in patients with recurrent self-harm. Single-centre double-blind randomised controlled trial. *Br J Psychiatry* 190: 118–122.
- Hautzinger M (2000). *Kognitive Verhaltenstherapie bei Depressionen*. 5. A. Weinheim: PVU/Beltz.
- Hayes SC, Follette VM, Linehan M (2004). *Mindfulness and Acceptance. Expanding the Cognitive-Behavioral Tradition*. New York: Guilford Press.
- Hecht H, van Calker D (2008). Evidenzbasierte Psychotherapie depressiver Störungen: Akuttherapie und Phasenprophylaxe. *Psychother Psych Med* 58: 1–13.

## 444.e4 11 Affektive Störungen

- Henkel V, Mergl R, Allgaier AK, et al. (2006). Treatment of depression with atypical features: a meta-analytic approach. *Psychiatry Res* 141(1): 89–101.
- Hollon SD, DeRubeis RJ, Shelton RC, et al. (2005). Prevention of relapse following cognitive therapy vs medications in moderate to severe depression. *Arch Gen Psychiatry* 62(4): 417.
- Hypericum Study Group (2002). Effect of *Hypericum perforatum* (St John's wort) in major depressive disorder: a randomized controlled trial. *JAMA* 287: 1807–1814.
- Imel ZE, Malterer MB, McKay KM, et al. (2008). A meta-analysis of psychotherapy and medication in unipolar depression and dysthymia. *J Affect Disord* 110: 197–206.
- Jakobsen JC, Lindschou Hansen J, Storebo OJ, et al. (2011). The effects of cognitive therapy versus 'treatment as usual' in patients with major depressive disorder. *PLoS One* 6(8): e22890.
- Jakobsen JC, Hansen JL, Simonsen S, et al. (2012). Effects of cognitive therapy versus interpersonal psychotherapy in patients with major depressive disorder: a systematic review of randomized clinical trials with meta-analyses and trial sequential analyses. *Psychol Med* 42(7): 1343–1357.
- Kanfer FH, Reinecker H, Schmelzer D (2006). *Selbstmanagementtherapie. Ein Lehrbuch für die klinische Praxis*. Berlin: Springer.
- Karyotaki E, Smit Y, Beurs DP, et al. (2016). The long-term efficacy of acute-phase psychotherapy for depression: a meta-analysis of randomized trials. *Depress Anxiety* 33(5): 370–383.
- Keck Jr PE, Marcus R, Tourkodimitris S, et al. for the Aripiprazole Study Group (2003a). A placebo-controlled, double-blind study of the efficacy and safety of aripiprazole in patients with acute bipolar mania. *Am J Psychiatry* 9: 1651–1658.
- Keck Jr PE, Versiani M, Potkin S, et al. (2003b) for the Ziprasidone in Mania Study Group: Ziprasidone in the treatment of acute bipolar mania: a three-week, placebo-controlled, double-blind, randomized trial. *Am J Psychiatry* 4: 741–748.
- Keck Jr PE, Calabrese JR, McQuade RD, et al. for the Aripiprazole Study Group (2006). A randomized, double-blind, placebo-controlled 26-week trial of aripiprazole in recently manic patients with bipolar I disorder. *J Clin Psychiatry* 4: 626–637.
- Keller MB, McCullough JP, Klein DN, et al. (2000). A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *N Engl J Med* 342: 1462–1470.
- Kim SW, Shin IS, Kim JM, et al. (2008). Effectiveness of mirtazapine for nausea and insomnia in cancer patients with depression. *Psychiatry Clin Neurosci* 1: 75–83.
- Klecha D, Legler M, Hüll M (2002). Zur Behandlung depressiver Störungen mit modernen Verfahren der Elektrokonvulsionstherapie. *Fortschr Neurol Psychiatr* 70: 353–367.
- Klein JP, Erkens N, Schweiger U, et al. (2018). Does childhood maltreatment moderate the effect of the Cognitive Behavioral Analysis System of Psychotherapy versus supportive psychotherapy in persistent depressive disorder? *Psychother Psychosom* 87(1): 46–48.
- Klerman GL, Weissman MM, Rounsville BJ, Chevron ES (1984). *Interpersonal Psychotherapy for Depression*. New York: Basic Books, pp. 71–182.
- Knekt P, Lindfors O, Häkkinen T, et al. (2008). Randomized trial on the effectiveness of long-and short-term psychodynamic psychotherapy and solution-focused therapy on psychiatric symptoms during a 3-year follow-up. *Psychol Med* 38(5): 689–703.
- Knekt P, Lindfors O, Sares-Jäské L, et al. (2013). Randomized trial on the effectiveness of long-and short-term psychotherapy on psychiatric symptoms and working ability during a 5-year follow-up. *Nord J Psychiatry* 67(1): 59–68.
- Köhler S, Wiethoff K, Ricken R, et al. (2015). Characteristics and differences in treatment outcome of inpatients with chronic vs. episodic major depressive disorders. *J Affect Disord* 173: 126–133.
- Krishna M, Jauhari A, Lepping P, et al. (2011). Is group psychotherapy effective in older adults with depression? A systematic review. *Int J Geriatr Psychiatry* 26(4): 331–340.
- Lam RW, Chan P, Wilkins-Ho M, Yatham LN (2008). Repetitive transcranial magnetic stimulation for treatment-resistant depression: a systematic review and metaanalysis. *Can J Psychiatry* 53: 621–631.
- Leichsenring F (2001). Comparative effects of short-term psychodynamic psychotherapy and cognitive-behavioral therapy in depression: a meta-analytic approach. *Clin Psychol Rev* 21(3): 401–419.
- Leichsenring F, Rabung S (2011). Long-term psychodynamic psychotherapy in complex mental disorders: update of a meta-analysis. *Br J Psychiatry* 199: 15–22.
- Leichsenring F, Steinert C (2017). Is cognitive behavioral therapy the gold standard for psychotherapy?: the need for plurality in treatment and research. *JAMA* 318(14): 1323–1324.
- Leichsenring F, Rabung S, Leibing E (2004). The efficacy of short-term psychodynamic psychotherapy in specific psychiatric disorders: a meta-analysis. *Arch Gen Psychiatry* 61(12): 1208–1216.
- Lewinsohn PM, Talkington J (1979). Studies on the measurement of unpleasant events and relations with depression. *Appl Psychol Meas* 3(1): 83–101.
- Lin PY, Su KP (2007). A meta-analytic review of double-blind, placebo-controlled trials of antidepressant efficacy of omega-3 fatty acids. *J Clin Psychiatry* 7: 1056–1061.
- Lisanby SH (2004). Magnetic seizure therapy: development of a novel convulsive technique. In: Lisanby SH (ed.). *Brain Stimulation in Psychiatric Treatment*. Washington, DC: American Psychiatric Publishing, pp. 67–98.
- Luoma J, Hayes SC, Walser RD (2009). ACT-Training: Handbuch der Acceptance & Commitment Therapie. Ein Lernprogramm in zehn Schritten. Paderborn: Junfermann.
- Ma SH, Teasdale JD (2004). Mindfulness-based cognitive therapy for depression: replication and exploration of differential relapse prevention effects. *J Consult Clin Psychol* 72: 31–40.
- Markowitz JC (2003). Interpersonal Psychotherapy for chronic depression. *J Clin Psychol* 59: 847–858.
- Mazzucchelli T, Kane R, Rees C (2009). Behavioral activation treatments for depression in adults: a meta-analysis and review. *Clin Psychol* 16(4): 383–411.
- McCullough JP (1991). Psychotherapy for dysthymia: a naturalistic study of ten patients. *J Nerv Ment Dis* 179: 734–740.
- McCullough JP (2000). *Treatment for Chronic Depression: Cognitive Behavioral Analysis System of Psychotherapy (CBASP)*. New York: Guilford Press.
- McDermut W, Miller IW, Brown RA (2001). The efficacy of group psychotherapy for depression: a meta-analysis and review of the empirical research. *Clin Psychol Sci Pract* 8: 98–116.
- McHugh RK, Whitton SW, Peckham AD, et al. (2013). Patient preference for psychological vs pharmacologic treatment of psychiatric disorders: a meta-analytic review. *J Clin Psychiatry* 74(6): 595–602.
- Meyer TD, Hautzinger M (2013). *Bipolare Störungen: Kognitiv-verhaltenstherapeutisches Behandlungsmanual*. Mit Online-Materialien. Weinheim: Beltz.
- Modell JG, Rosenthal NE, Harriett AE, et al. (2005). Seasonal affective disorder and its prevention by anticipatory treatment with bupropion XL. *Biol Psychiatry* 58(8): 658–667.
- Negt P, Brakemeier EL, Michalak J, et al. (2016). The treatment of chronic depression with cognitive behavioral analysis system of psychotherapy: a systematic review and meta-analysis of randomized controlled clinical trials. *Brain Behav* 6(8): 2162–3279.
- Nelson JC (1999). A review of the efficacy of serotonergic and noradrenergic reuptake inhibitors for treatment of major depression. *Biol Psychiatry* 46: 1301–1308.
- Nemeroff CB, Heim CM, Thase ME, et al. (2003). Differential responses to psychotherapy versus pharmacotherapy in patients with chronic forms of major depression and childhood trauma. *Proc Natl Acad Sci USA* 100: 14293–14296.
- Nemets H, Nemets B, Apter A, et al. (2006). Omega-3 treatment of childhood depression: a controlled, double-blind pilot study. *Am J Psychiatry* 163: 1098–1100.

- Nieuwstraten CE, Dolovich LR (2001). Bupropion versus selective serotonin-reuptake inhibitors for treatment of depression. *Ann Pharmacother* 35: 1608–1613.
- O'Reardon JP, Solvason B, Janicak PG, et al. (2007). Efficacy and safety of transcranial magnetic stimulation in the acute treatment of major depression: a multi-site randomized controlled trial. *Biol Psychiatry* 15: 553–563.
- Owen C, Rees AM, Parker G (2008). The role of fatty acids in the development and treatment of mood disorders. *Curr Opin Psychiatry* 1: 19–24.
- Piaget J (1923/1926). *The Language and Thought of the Child*. New York: Harcourt Brace.
- Poolsup N, Li Wan Po A, de Oliveira IR (2000). Systematic overview of lithium treatment in acute mania. *J Clin Pharm Ther* 25: 139–156.
- Potkin SG, Keck PE Jr, Segal S, et al. (2005). Ziprasidone in acute bipolar mania: a 21-day randomized, double-blind, placebo-controlled replication trial. *J Clin Psychopharmacol* 4: 301–310.
- Ravindran AV, Anisman H, Merali Z, et al. (1999). Treatment of primary dysthymia with group cognitive therapy and pharmacotherapy: clinical symptoms and functional impairments. *Am J Psychiatry* 156: 1608–1617.
- Reynolds CF, Kupfer DJ (1999). Depression and aging: a look to the future. *Psychiatr Serv* 50: 1167–1172.
- Rush AJ, Marangell LB, Sackeim HA, et al. (2005). Vagus nerve stimulation for treatment-resistant depression: a randomized, controlled acute phase trial. *Biol Psychiatry* 58: 347–354.
- Sachs GS, Lafer B, Stoll AL, et al. (1994). A double-blind trial of bupropion versus desipramine for bipolar depression. *J Clin Psychiatry* 55: 391–393.
- Sackeim HA, Prudic J, Devanand DP, et al. (2000). Prospective, randomized, double-blind comparison of bilateral and right unilateral electroconvulsive therapy at different stimulus intensities. *Arch Gen Psychiatry* 57: 425–434.
- Sackeim HA, Prudic J, Nobler MS, et al. (2008). Effects of pulse width and electrode placement on the efficacy and cognitive effects of electroconvulsive therapy. *Brain Stimul* 1(2): 71–83.
- Samad Z, Brealey S, Gilbody S (2011). The effectiveness of behavioural therapy for the treatment of depression in older adults: a meta-analysis. *Int J Geriatr Psychiatry* 26(12): 1211–1220.
- Schramm E, Berger M (2012). Replik zu „Psychodynamische Therapie der Depression“ von Leichsenring F, Rabung S. *Nervenarzt*: DOI 10.1007/s00115-012-3545-2 (<http://link.springer.com/article/10.1007/s00115-012-3545-2#page-1>).
- Schramm E, Klecha D (2010). *Interpersonelle Psychotherapie in der Gruppe. Das Kurzmanual*. Stuttgart: Schattauer.
- Schramm E, van Calker D, Dykier P, et al. (2007). An intensive treatment program of Interpersonal Psychotherapy plus pharmacotherapy for depressed inpatients: acute and long-term results. *Am J Psychiatry* 164: 768–777.
- Schramm E, Schneider D, Zobel I, et al. (2008). Efficacy of Interpersonal Psychotherapy plus pharmacotherapy in chronically depressed inpatients. *J Affect Disord* 9(1–2): 65–73.
- Schramm E, Zobel I, Dykier P, et al. (2010). Cognitive Behavioral Analysis System of Psychotherapy versus Interpersonal Psychotherapy for early-onset chronic depression: a randomized pilot study. *J Affect Disord* 129(1–3): 109–116.
- Schramm E, Kriston L, Zobel I, et al. (2017). Effect of disorder-specific vs nonspecific psychotherapy for chronic depression: a randomized clinical trial. *JAMA Psychiatry* 74(3): 233–242.
- Segal ZV, Williams JMG, Teasdale JD (2002). *Mindfulness-Based Cognitive Therapy for Depression: A New Approach to Preventing Relapse*. New York: Guilford Press.
- Seligman MEP (1975). *Helplessness*. San Francisco: Freeman.
- Shear K, Frank E, Houck PR, Reynolds CF (2005). Treatment of complicated grief: a randomized controlled trial. *JAMA* 293: 2658–2660.
- Small JG, Klapper MH, Kellams JJ, et al. (1998). Electroconvulsive treatment compared with lithium in the management of manic states. *Arch Gen Psychiatry* 45: 727–732.
- Smith D, Dempster C, Glanville J, et al. (2002). Efficacy and tolerability of venlafaxin compared with selective serotonin reuptake inhibitors and other antidepressants: a meta-analysis. *Br J Psychiatry* 180: 396–404.
- Spaans HP, Kho KH, Verwijk E, et al. (2013). Efficacy of ultrabrief pulse electroconvulsive therapy for depression: a systematic review. *J Affect Disord* 150(3): 720–726.
- Stassen HH, Angst J, Hell D, et al. (2007). Is there a common resilience mechanism underlying antidepressant drug response? Evidence from 2,848 patients. *J Clin Psychiatry* 68: 1195–1205.
- Stenzel N, Bollmann S, Brakemeier EL (2017). Depression. In: Brakemeier EL, Jacobi F (Hrsg.). *Verhaltenstherapie in der Praxis*. Weinheim: Beltz, S. 500–512.
- Strauss C, Cavanagh K, Oliver A, Pettman D (2014). Mindfulness-based interventions for people diagnosed with a current episode of an anxiety or depressive disorder: a meta-analysis of randomised controlled trials. *PLOS One* 9(4): e96110.
- Stulz N, Thase ME, Klein DN, et al. (2010). Differential effects of treatments for chronic depression: a latent growth model reanalysis. *J Consult Clin Psychol* 78(3): 409.
- Thase ME, Greenhouse JB, Frank E, et al. (1997). Treatment of major depression with psychotherapy or psychotherapy-pharmacotherapy combinations. *Arch Gen Psychiatry* 54(11): 1009–1015.
- Thase ME, Macfadden W, Weisler RH, et al. for the BOLDER II Study Group (2006). Efficacy of quetiapine monotherapy in bipolar I and II depression: a double-blind, placebo-controlled study (the BOLDER II study). *J Clin Psychopharmacol* 26(6): 600–609.
- Tolin DF (2010). Is cognitive-behavioral therapy more effective than other therapies? A meta-analytic review. *Clin Psychol Rev* 30: 710–720.
- Tondo L, Hennen J, Baldessarini RJ (2003). Rapid-cycling bipolar disorder: effects of long-term treatments. *Acta Psychiatr Scand* 108(1): 4–14.
- UK ECT Review Group (2003). Efficacy and safety of electroconvulsive therapy in depressive disorders: a systematic review and meta-analysis. *Lancet* 361: 799–808.
- Vieta E, Mullen J, Brecher M, et al. (2005). Quetiapine monotherapy for mania associated with bipolar disorder: combined analysis of two international, double-blind, randomised, placebo-controlled studies. *Curr Med Res Opin* 6: 923–934.
- Vis PM, van Baardewijk M, Einarson TR (2005). Duloxetin and venlafaxin-XR in the treatment of major depressive disorder: a meta-analysis of randomized clinical trials. *Ann Pharmacother* 39: 1798–1807.
- Wani AL, Bhat SA, Ara A (2015). Omega-3 fatty acids and the treatment of depression: a review of scientific evidence. *Integr Med Res* 3: 132–141.
- Watson JC, Bedard DL (2006). Client's emotional processing in psychotherapy: a comparison between cognitive-behavioral and process-experiential therapies. *J Consult Clin Psychol* 74: 152–159.
- Watson JC, Gordon LB, Stermac L, et al. (2003). Comparing the effectiveness of process-experiential with cognitive behavioral psychotherapy in the treatment of depression. *J Consult Clin Psychol* 71: 773–781.
- Weiner RD, Rogers HJ, Davidson JR, Kahn EM (1986). Effects of electroconvulsive therapy upon brain electrical activity. *Ann N Y Acad Sci* 462(1): 270–281.
- Weissman MM, Markowitz JC, Klerman G (2017). *The Guide to Interpersonal Psychotherapy: Updated and expanded*. New York: Oxford University Press.
- Wenzel-Seifert K, Wittmann M, Haen E (2010). Torsade de pointes episodes under treatment with selective serotonin reuptake inhibitors. *Pharmacopsychiatry* 43(7): 279–281.
- Wiersma, JE, Hovens, JGFM, van Oppen P, et al. (2009). The importance of early trauma and childhood life events for chronicity of depression. *J Clin Psychiatry* 70: 983–989.
- Wittchen H-U, Möller HJ, Vossen A et al. (1995). Depression. *Wege aus der Krankheit*. Basel, Freiburg, Berlin: Karger.
- WHO – World Health Organization (2017). *Global Strategy for Women's, Children's and Adolescents' Health (2016–2030)*. Geneva: World Health Organization.
- von Wolff A, Hözel LP, Westphal A, et al. (2012). Combination of pharmacotherapy and psychotherapy in the treatment of chronic depression: a systematic review and meta-analysis. *BMC Psychiatry* 12(1): 61.

- Wu JC, Gillin JC, Buchsbaum MS, et al. (1992). Effect of sleep deprivation on brain metabolism of depressed patients. *Am J Psychiatry* 149: 538–543.
- Zhou X, Hetrick SE, Cuijpers P, et al. (2015). Comparative efficacy and acceptability of psychotherapies for depression in children and adolescents: a systematic review and network meta-analysis. *World Psychiatry* 14: 207–222.
- 11.7 ERHALTUNGSTHERAPIE UND REZIDIVPROPHYLAXE**
- Altshuler L, Suppes T, Black D, et al. (2003). Impact of antidepressant discontinuation after acute bipolar depression remission on rates of depressive relapse at 1-year follow-up. *Am J Psychiatry* 7: 1252–1262.
- APA Committee on Electroconvulsive Therapy (2001). The Practice of Electroconvulsive Therapy: Recommendations for Treatment, Training and Privileging. 2<sup>nd</sup> ed. Washington, DC: American Psychiatric Association.
- Beynon S, Soares-Weiser K, Woolacott N, et al. (2008). Psychosocial interventions for the prevention of relapse in bipolar disorder: systematic review of controlled trials. *Br J Psychiatry* 1: 5–11.
- Brakemeier EL, Merkl A, Wilbertz G, et al. (2013). Cognitive-behavioral therapy as continuation treatment to sustain response after electroconvulsive therapy in depression: a randomized controlled trial. *Biol Psychiatry* 76(3): 194–202.
- Dardennes R, Even C, Bange F, Heim A (1995). Comparison of carbamazepine and lithium in the prophylaxis of bipolar disorders. A meta-analysis. *Br J Psychiatry* 166(3): 378–381.
- De Mello FM, de Jesus M, Bacalchuk J, et al. (2005). A systematic review of research findings on the efficacy of interpersonal therapy for depressive disorders. *Eur Arch Psychiatry Clin Neurosci* 255(2): 75–82.
- Fava GA, Tomba E, Grandi S (2007). The road to recovery from depression – don't drive today with yesterday's map. *Psychother Psychosom* 76(5): 260–265.
- Frank E (2007). Interpersonal and social rhythm therapy: a means of improving depression and preventing relapse in bipolar disorder. *J Clin Psychol* 5: 463–473.
- Frank E, Johnson S, Kupfer DJ (1992). Psychological treatments in prevention of relapse. In: Montgomery SA, Rouillon F (eds.). Long-Term Treatment of Depression. New York: Wiley, pp. 197–228.
- Freeman MP, Fava M, Lake J, et al. (2010). Complementary and alternative medicine in major depressive disorder: the American Psychiatric Association Task Force report. *J Clin Psychiatry* 71(6): 669.
- Geschwind N, Peeters F, Hubers M, et al. (2012). Efficacy of mindfulness-based cognitive therapy in relation to prior history of depression: randomised controlled trial. *Br J Psychiatry* 2014(4): 320–325.
- Goodwin FK, Jamison KR (1990). Manic-Depressive Illness. Oxford, New York, Tokyo: Oxford University Press.
- Greil W, Ludwig-Mayerhofer W, Erazo N, et al. (1996). Comparative efficacy of lithium and amitriptyline in the maintenance treatment of recurrent unipolar depression: a randomised study. *J Affect Disord* 40: 179–190.
- Greil W, Ludwig-Mayerhofer W, Erazo N, et al. (1997). Lithium versus carbamazepine in the maintenance treatment of bipolar disorders – a randomised study. *J Affect Disord* 43: 151–161.
- Greil W, Kleindienst N, Erazo N, Müller-Oerlinghausen B (1998). Differential response to lithium and carbamazepine in the prophylaxis of bipolar disorder. *J Clin Psychopharmacol* 18: 455–460.
- Greil W, Kleindienst N (1999a). The comparative prophylactic efficacy of lithium and carbamazepine in patients with bipolar I disorder. *Int Clin Psychopharmacol* 14: 277–281.
- Greil W, Kleindienst N (1999b). Lithium versus carbamazepine in the maintenance treatment of bipolar II disorder and bipolar disorder not otherwise specified. *Int Clin Psychopharmacol* 14: 283–285.
- Grünfeld JP, Rossier BC (2009). Lithium nephrotoxicity revisited. *Nat Rev Nephrol* 5: 270–276.
- Guzzetta F, Tondo L, Centorrini F, Baldessarini RJ (2007). Lithium treatment reduces risk in recurrent major depressive disorder. *J Clin Psychiatry* 68: 380–383.
- Henkel V, Mergl R, Allgaier AK, et al. (2006). Treatment of depression with atypical features: a meta-analytic approach. *Psychiatry Res* 141: 89–101.
- Honig A, Arts BM, Ponds RW, Riedel WJ (1999). Lithium-induced cognitive side-effects in bipolar disorder: a qualitative analysis and implications for daily practice. *Int Clin Psychopharmacol* 14: 167–171.
- Lam DH, Watkins ER, Hayward P, et al. (2003). A randomised controlled study of cognitive therapy for relapse prevention for bipolar affective disorder. *Arch Gen Psychiatry* 60: 145–152.
- Ma SH, Teasdale JD (2004). Mindfulness-based cognitive therapy for depression: replication and exploration of differential relapse prevention effects. *J Consult Clin Psychol* 72: 31–40.
- Montgomery SA, Rasmussen JG, Tanghoj P (1993). A 24-week study of 20 mg citalopram, 40 mg citalopram, and placebo in the prevention of relapse of major depression. *Int Clin Psychopharmacol* 8: 181–188.
- Oestergaard S, Møldrup C (2011). Optimal duration of combined psychotherapy and pharmacotherapy for patients with moderate and severe depression: a meta-analysis. *J Affect Disord* 131(1–3): 24–36.
- Paykel ES (2007). Cognitive therapy in relapse prevention in depression. *Int J Neuropsychopharmacol* 10: 131–136.
- Teasdale JD, Segal ZV, Williams JM, et al. (2000). Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *J Consult Clin Psychol* 68: 615–623.
- Tondo L, Hennen J, Baldessarini RJ (2003). Rapid-cycling bipolar disorder: effects of long-term treatments. *Acta Psychiatr Scand* 108: 4–14.
- Vittengl JR, Clark LA, Dunn TW, Jarrett RB (2007). Reducing relapse and recurrence in unipolar depression: a comparative meta-analysis of cognitive-behavioral therapy's effects. *J Consult Clin Psychol* 75: 475–488.
- Yatham LN, Kusumakar V, Calabrese JR, et al. (2002). Third generation anticonvulsants in bipolar disorder: a review of efficacy and summary of clinical recommendations. *J Clin Psychiatry* 63: 275–283.
- LEITLINIEN (SYSTEMATISCH EVIDENZBASIERT)**
- Anderson IM, Ferrier IN, Baldwin RC, et al. (2008). Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2000 British Association for Psychopharmacology guidelines. *J Psychopharmacol* 22: 343–396.
- APA – American Psychiatric Association (2002). Practice Guidelines for the Treatment of Patients with Bipolar Disorder. 2<sup>nd</sup> ed. Washington, DC: American Psychiatric Press.
- APA – American Psychiatric Association (2010). Practice Guideline for the Treatment of Patients with Major Depressive Disorder; [www.psychiatryonline.com/pracGuide/pracGuideTopic\\_7.aspx](http://www.psychiatryonline.com/pracGuide/pracGuideTopic_7.aspx) (letzter Zugriff: 26.6.2018).
- Deutsche Gesellschaft für Bipolare Störung (DGBS), Deutsche Gesellschaft für Psychiatrie, Psychotherapie und Nervenheilkunde (DGPPN) (2011). S3-Leitlinie zur Diagnostik und Therapie bipolarer Störungen; [www.leitlinie-bipolar.de/](http://www.leitlinie-bipolar.de/).
- Deutsche Gesellschaft für Psychiatrie, Psychotherapie und Nervenheilkunde (DGPPN), Bundesärztekammer (BÄK), Kassenärztliche Bundesvereinigung (KBV) und Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF) (Hrsg.) (2010). Nationale VersorgungsLeitlinie Unipolare Depression. S3-Leitlinien in Psychiatrie und Psychotherapie. Berlin: Springer.
- DGPPN, BÄK, KBV AWMF (2015). S3-Leitlinie/Nationale VersorgungsLeitlinie Unipolare Depression – Langfassung. 2. Auflage. Version 5; [www.depression.versorgungsleitlinien.de](http://www.depression.versorgungsleitlinien.de) (letzter Zugriff: 15.12.2017).
- Härter M, Bermejo I, Dausend S, et al. (2014). VersorgungsLeitlinien für depressive Störungen in der ambulanten Praxis. ZEFQ: Supplementband.
- Lam RW, Kennedy SH, Grigoriadis S, et al. (2009). Canadian Network for Mood and Anxiety Treatments (CANMAT) Clinical guidelines for the management of major depressive disorder in adults. II. Pharmacotherapy. *J Affect Disord* 117: S26–S43.
- National Institute for Clinical Excellence (NICE) (2006). Bipolar disorder. The Management of Bipolar Disorder in Adults, Children and Adolescents, in Primary and Secondary Care. Clinical Guideline 38. London, NICE: <http://guidance.nice.org.uk/CG38/Guidance/pdf/English> (letzter Zugriff: 26.6.2018).
- National Institute for Clinical Excellence (NICE) (2010). Depression: Management of Depression in Primary and Secondary Care. Clinical Guideline 90.

- London, NICE: <http://guidance.nice.org.uk/CG90/Guidance/pdf/English> (letzter Zugriff: 26.6.2018).
- Parik SV, Segal ZV, Grigoriadis S, et al. (2009). Canadian Network for Mood and Anxiety Treatments (CANMAT) Clinical guidelines for the management of major depressive disorder in adults. II. Psychotherapy alone or in combination with antidepressant medication. *J Affect Disord* 117: S15–S25.
- Suehs BT, Argo TR, Bendele SD, et al. (2008). Texas Medication Algorithm Project Procedural Manual: Major Depressive Disorder Algorithms. Austin, TX, Texas Department of State Health Services.

#### **SYSTEMATISCHE COCHRANE-REVIEWS (WWW.COCHRANE.DE/DEUTSCH)**

- Abbas AA, Kisely SR, Town JR, et al. (2014). Short-term psychodynamic psychotherapies for common mental disorders. *Cochrane Database Syst Rev* 7: CD004687.
- Appleton KM, Sallis HM, Perry R, et al. (2015). Omega-3 fatty acids for depression in adults. *Cochrane Database Syst Rev* 11: CD004692.
- Arroll B, Elley CR, Fishman T, et al. (2009). Antidepressants versus placebo for depression in primary care. *Cochrane Database Syst Rev* 3: CD007954.
- Barbato A, D'Avanzo B (2006). Marital therapy for depression. *Cochrane Database Syst Rev* 2: CD004188.
- Baumeister H, Hutter N, Bengel J (2011). Psychological and pharmacological interventions for depression in patients with coronary artery disease. *Cochrane Database Syst Rev* 2: CD008012.
- Brown R, Taylor MJ, Geddes J (2013). Aripiprazole alone or in combination for acute mania. *Cochrane Database Syst Rev* 12: CD005000.
- Burgess S, Geddes JR, Hawton K, et al. (2001). Lithium for maintenance treatment of mood disorders. *Cochrane Database Syst Rev* 3: CD003013.
- Caddy C, Amit BH, McCloud TL, et al. (2015). Ketamine and other glutamate receptor modulators for depression in adults. *Cochrane Database Syst Rev* 9: CD011612.
- Cipriani A, Rendell JM, Geddes JR (2006a). Haloperidol alone or in combination for acute mania. *Cochrane Database Syst Rev* 3: CD004362.
- Cipriani A, Smith K, Burgess S, et al. (2006b). Lithium versus antidepressants in the long-term treatment of unipolar affective disorder. *Cochrane Database Syst Rev* 4: CD003492.
- Cipriani A, Purgato M, Furukawa TA, et al. (2012a). Citalopram versus other anti-depressive agents for depression. *Cochrane Database Syst Rev* 7: CD006534.
- Cipriani A, Koesters M, Furukawa TA, et al. (2012b). Duloxetine versus other anti-depressive agents for depression. *Cochrane Database Syst Rev* 10: CD006533.
- Cipriani A, Reid K, Young AH, et al. (2013). Valproic acid, valproate and divalproex in the maintenance treatment of bipolar disorder. *Cochrane Database Syst Rev* 10: CD003196.
- Dennis C-L, Hodnett E (2007). Psychosocial and psychological interventions for treating postpartum depression. *Cochrane Database Syst Rev* 4: CD006116.
- Eshun-Wilson I, Siegfried N, Akena DH (2018). Antidepressants for depression in adults with HIV infection. *Cochrane Database Syst Rev* 1: CD008525.
- Forneris CA, Nussbaumer B, Kaminski-Hartenthaler A, et al. (2015). Psychological therapies for preventing seasonal affective disorder. *Cochrane Database Syst Rev* 11: CD011270.
- Furukawa TA, Streiner DL, Young LT, Kinoshita Y (2001). Antidepressant plus benzodiazepine for major depression. *Cochrane Database Syst Rev* 2: CD001026.
- Furukawa T, McGuire H, Barbui C (2003). Low dosage tricyclic antidepressants for depression. *Cochrane Database Syst Rev* 3: CD003197.
- Gartlehner G, Nussbaumer B, Gaynes BN, et al. (2015). Second-generation antidepressants for preventing seasonal affective disorder in adults. *Cochrane Database Syst Rev* 11: CD011268.
- Guajana G, Barbui C, Hotopf M (2007). Amitriptyline for depression. *Cochrane Database Syst Rev* 2: CD004186.
- Guajana G, Gupta S, Chiodo D, et al. (2013). Agomelatine versus other anti-depressive agents for major depression. *Cochrane Database Syst Rev* 12: CD008851.
- Hackett ML, Anderson CS, House A, Xia J (2008). Interventions for treating depression after stroke. *Cochrane Database Syst Rev* 4: CD003437.
- Henken HT, Huibers MJH, Churchill R, et al. (2007). Family therapy for depression. *Cochrane Database Syst Rev* 3: CD0006728.
- Hoffbrand S, Howard L, Crawley H (2001). Antidepressant treatment for post-natal depression. *Cochrane Database Syst Rev* 2: CD0002018.
- Howard LM, Hoffbrand S, Henshaw C, et al. (2005). Antidepressant prevention of postnatal depression. *Cochrane Database Syst Rev* 2: CD004363.
- Hunot V, Moore THM, Caldwell DM, et al. (2013). 'Third wave' cognitive and behavioural therapies versus other psychological therapies for depression. *Cochrane Database Syst Rev* 10: CD008704.
- Kaminski-Hartenthaler A, Nussbaumer B, Forneris CA, et al. (2015). Melatonin and agomelatine for preventing seasonal affective disorder. *Cochrane Database Syst Rev* 11: CD011271.
- Koesters M, Ostuzzi G, Guajana G, et al. (2017). Vortioxetine for depression in adults. *Cochrane Database Syst Rev* 7: CD011520. DOI: <https://doi.org/10.1002/14651858.CD011520>.
- Leucht C, Huhn M, Leucht S (2012). Amitriptyline versus placebo for major depressive disorder. *Cochrane Database Syst Rev* 12: CD009138.
- Linde K, Berner MM, Kriston L (2008). St John's wort for major depression. *Cochrane Database Syst Rev* 4: CD000448.
- Macritchie K, Geddes JR, Scott J, et al. (2003a). Valproate for acute mood episodes in bipolar disorder. *Cochrane Database Syst Rev* 1: CD004052.
- Macritchie KAN, Geddes JR, Scott J, et al. (2003b). Valproic acid, valproate and divalproex in the maintenance treatment of bipolar disorder. *Cochrane Database Syst Rev* 3: CD003196.
- Magni LR, Purgato M, Gastaldon C, et al. (2013). Fluoxetine versus other types of pharmacotherapy for depression. *Cochrane Database Syst Rev* 7: CD004185.
- McCloud TL, Caddy C, Jochim J (2015). Ketamine and other glutamate receptor modulators for depression in bipolar disorder in adults. *Cochrane Database Syst Rev* 9: CD011611.
- Morriss RK, Faizal MA, Jones AP, et al. (2007). Interventions for helping people recognise early signs of recurrence in bipolar disorder. *Cochrane Database Syst Rev* 1: CD004854.
- Mottram P, Wilson K, Strobl J (2006). Antidepressants for depressed elderly. *Cochrane Database Syst Rev* 1: CD003491.
- Nussbaumer B, Kaminski-Hartenthaler A, Forneris CA, et al. (2015). Light therapy for preventing seasonal affective disorder. *Cochrane Database Syst Rev* 11: CD011269.
- Ostuzzi G, Matcham F, Dauchy S, et al. (2015). Antidepressants for the treatment of depression in people with cancer. *Cochrane Database Syst Rev* 6: CD011006.
- Pigott K, Galizia I, Vasudev K, et al. (2016). Topiramate for acute affective episodes in bipolar disorder in adults. *Cochrane Database Syst Rev* 9: CD003384.
- Rayner L, Price A, Evans A, et al. (2010). Antidepressants for depression in physically ill people. *Cochrane Database Syst Rev* 3: CD007503.
- Rendell JM, Gijsman HJ, Keck P, et al. (2003). Olanzapine alone or in combination for acute mania. *Cochrane Database Syst Rev* 3: CD004040.
- Rendell JM, Gijsman HJ, Bauer MS, et al. (2006). Risperidone alone or in combination for acute mania. *Cochrane Database Syst Rev* 1: CD004043.
- Rodriguez-Martin JL, Barbanjo JM, Schlaepfer TE, et al. (2001). Transcranial magnetic stimulation for treating depression. *Cochrane Database Syst Rev* 4: CD003493.
- Shinohara K, Honyashiki M, Imai H, et al. (2013). Behavioural therapies versus other psychological therapies for depression. *Cochrane Database Syst Rev* 10: CD008696.
- Silva de Lima M, Hotopf M. (2003). Pharmacotherapy for dysthymia. *Cochrane Database Syst Rev* 3: CD004047.
- Stek ML, van der Wurff FFB, Hoogendoijk WJG, Beekman ATF (2003). Electroconvulsive therapy for the depressed elderly. *Cochrane Database Syst Rev* 2: CD003593.

## 444.e8 11 Affektive Störungen

- Taylor MJ, Rudkin L, Bullemer-Day P, et al. (2013). Strategies for managing sexual dysfunction induced by antidepressant medication. Cochrane Database Syst Rev 5: CD003382.
- Thaler K, Delivuk M, Chapman A, et al. (2011). Second-generation antidepressants for seasonal affective disorder. Cochrane Database Syst Rev 12: CD008591.
- Tuunainen A, Kripke DF, Endo T (2004). Light therapy for non-seasonal depression. Cochrane Database Syst Rev 2: CD004050.
- Watanabe N, Omori IM, Nakagawa A, et al. (2011). Mirtazapine versus other antidepressive agents for depression. Cochrane Database Syst Rev 12: CD006528.
- Wijkstra J, Lijmer J, Burger H et al. (2015). Pharmacological treatment for psychotic depression. Cochrane Database Syst Rev 7: CD004044.
- Wilkinson P, Izmeth Z (2016). Continuation and maintenance treatments for depression in older people. Cochrane Database Syst Rev 9: CD006727.
- Wilson K, Mottram P, Sivanranthan A, Nightingale A (2001). Antidepressants versus placebo for the depressed elderly. Cochrane Database Syst Rev 1: CD000561.
- Wilson KCM, Mottram PG, Vassilas CA (2008). Psychotherapeutic treatments for older depressed people. Cochrane Database Syst Rev 1: CD004853.

### PATIENTENRATGEBER

- Hautzinger M (2006). Ratgeber Depression. Informationen für Betroffene und Angehörige. Göttingen, Bern, Toronto: Hogrefe.
- Meyer TD, Hautzinger M (2013). Ratgeber Manisch-depressive Erkrankungen. Informationen für Menschen mit einer bipolaren Störung und deren Angehörige. Göttingen: Hogrefe.
- Hegerl U, Niescken S (2013). Depressionen bewältigen, die Lebensfreude wiederfinden. Stuttgart: Trias.
- Müller-Rörich T, Hass K, Margue F (2013). Schattendasein: Das unverstandene Leiden Depression. Berlin, Heidelberg: Springer.