

LITERATUR

19.1 GRUNDLAGEN DER SCHLAFFORSCHUNG

- Aserinsky E, Kleitman N (1953). Regularly occurring periods of eye motility and concomitant phenomena during sleep. *Science* 118: 273–274.
- Carskadon MA, Dement WC, Mitler MM, et al. (1986). Guidelines for the multiple sleep latency test (MSLT). A standard measure of sleepiness. *Sleep* 9: 519–524.
- Iber C, Ancoli-Israel S, Chesson A, et al. (2007). The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications. Westchester, Illinois: American Academy of Sleep Medicine.
- Kryger MH, Roth T, Dement WC (eds.) (2011). Principles and Practice of Sleep Medicine. 5th ed. Philadelphia: Saunders.
- Littner MR, Kushida C, Wise M, et al. (2005). Standards of Practice Committee of the American Academy of Sleep Medicine. Practice parameters for clinical use of the multiple sleep latency test and the maintenance of wakefulness test. *Sleep* 28: 113–121.
- Rechtschaffen A, Kales A (1968). A Manual of Standardized Terminology, Techniques and Scoring System for Sleep Stages of Human Subjects. Washington DC: U. S. Government Printing Office.
- Redline S, Kirchner HL, Quan SF, et al. (2004). The effects of age, sex, ethnicity, and sleep-disordered breathing on sleep architecture. *Arch Intern Med* 164: 406–418.

19.2 KLASSEKATION DER SCHLAFSTÖRUNGEN

- AASM – American Academy of Sleep Medicine (2014). The International Classification of Sleep Disorders. 3rd ed. Darien, IL: AASM.
- APA – American Psychiatric Association (2013). Diagnostic and Statistical Manual of Mental Disorders, 5th edition DSM-5™. Washington, DC: APA.
- APA – American Psychiatric Association (2015). Diagnostisches und Statistisches Manual Psychischer Störungen DSM-5. Deutsche Ausgabe hrsg. von P. Falkai und H-U Wittchen. Göttingen, Bern, Wien: Hogrefe.
- Dilling H, Mombour W, Schmidt MH (1991). Internationale Klassifikation psychischer Störungen: ICD-10 Kapitel V (F), Klinisch-diagnostische Leitlinien. 1. A. Bern, Göttingen, Toronto: Huber.
- Mayer G, Fietze I, Fischer J et al. (2009). Nicht erholsamer Schlaf. Leitlinie S3 der DGSM. Wien: Blackwell.
- Schramm E, Riemann D (Hrsg.) (1995). Internationale Klassifikation der Schlafstörungen. Weinheim: PVU.

19.3 PRIMÄRE SCHLAFSTÖRUNGEN

- Alajmi M, Mulgrew AT, Fox J, et al. (2007). Impact of continuous positive airway pressure therapy on blood pressure in patients with obstructive sleep apnea hypopnea: a meta-analysis of randomized controlled trials. *Lung* 185: 67–72.
- Backhaus J, Hohagen F, Voderholzer U, Riemann D (2001). Long-term effectiveness of a short-term cognitive-behavioral group treatment for primary insomnia. *Eur Arch Psychiatry Clin Neurosci* 251: 35–41.
- Baglioni C, Battagliese G, Feige B, et al. (2011). Insomnia as a clinical predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord* 135: 10–19.
- Bastien CH, Vallieres A, Morin CM (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med* 2: 297–307.
- Billiard M, Dauvilliers Y (2001). Idiopathic hypersomnia. *Sleep Med Rev* 5: 349–358.
- Breslau N, Roth T, Rosenthal L, Andreski P (1996). Sleep disturbance and psychiatric disorders: a longitudinal epidemiological study of young adults. *Biol Psychiatry* 39: 411–418.
- Broughton RJ (1968). Sleep disorders: disorders of arousal? Enuresis, somnambulism, and nightmares occur in confusional states of arousal, not in "dreaming sleep". *Science* 159: 1070–1078.
- Buscemi N, Vandermeer B, Hooton N, et al. (2005). The efficacy and safety of exogenous melatonin for primary sleep disorders. A meta-analysis. *J Gen Intern Med* 20: 1151–1158.
- Buyse DJ, Reynolds CF 3rd, Monk TH, et al. (1989). The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 28: 193–213.
- Cheng SK, Dizon J (2012). Computerised cognitive behavioural therapy for insomnia: a systematic review and meta-analysis. *Psychother Psychosom* 81: 206–216.
- Chesson AL Jr, Ferber RA, Fry JM, et al. (1997). The indications for polysomnography and related procedures. *Sleep* 20: 423–487.
- Dreßing H, Riemann D (1994). Diagnostik und Therapie von Schlafstörungen. Stuttgart, Jena: Fischer.
- Glass J, Lancot KL, Herrmann N, et al. (2005). Sedative hypnotics in older people with insomnia: meta-analysis of risks and benefits. *BMJ* 331: 1169.
- Hohagen F, Rink K, Schramm E, et al. (1993). Prevalence and treatment of insomnia in general practice – a longitudinal study. *Eur Arch Psy Clin N* 242: 329–336.
- Huedo-Medina TB, Kirsch I, Mittlemass J, et al. (2012). Effectiveness of non-benzodiazepine hypnotics in treatment of adult insomnia: meta-analysis of data submitted to the Food and Drug Administration. *BMJ* 356: e8343.
- Irwin MR, Cole JC, Nicassio PM (2006). Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+ years of age. *Health Psychol* 25: 3–14.
- Koffel EA, Koffel JB, Gehrman PR (2015). A meta-analysis of group cognitive behavioral therapy for insomnia. *Sleep Med Rev* 19: 6–16.
- Krystal AD, Lankford A, Durrence HH, et al. (2011). Efficacy and safety of doxepin 3 and 6 mg in a 35-day sleep laboratory trial in adults with chronic primary insomnia. *Sleep* 34: 1433–1442.
- Leach MJ, Page AT (2015). Herbal medicine for insomnia: a systematic review and meta-analysis. *Sleep Med Rev* 24: 1–12.
- Li M, Zhang XW, Hou WS, Tang ZY (2014). Insomnia and risk of cardiovascular disease: a meta-analysis of cohort studies. *Int J Cardiol* 176: 1044–1047.
- Littner M, Johnson SF, McCall WV, et al. (2001). Practice parameters for the treatment of narcolepsy: an update for 2000. *Sleep* 24: 451–466.
- McMahon JP, Foresman BH, Chisholm RC (2003). The influence of CPAP on the neurobehavioral performance of patients with obstructive sleep apnea hypopnea syndrome: a systematic review. *WMJ* 102: 36–43.
- Morin CM (1993). Insomnia. New York, London: Guilford Press.
- Morin CM, Culbert JP, Schwartz SM (1994). Nonpharmacological interventions for insomnia: a meta-analysis of treatment efficacy. *Am J Psychiatry* 151: 1172–1180.
- Morin CM, Colecci C, Stone J, et al. (1999). Behavioral and pharmacological therapies for late-life insomnia: a randomized controlled trial. *JAMA* 281: 991–999.
- Morin CM, Vallieres A, Guay B, et al. (2009). Cognitive behavioral therapy, singly and combined with medication, for persistent insomnia. *JAMA* 301: 2005–2015.
- Murtagh DR, Greenwood KM (1995). Identifying effective psychological treatments for insomnia: a meta-analysis. *J Consult Clin Psychol* 63: 79–89.
- Nowell PD, Mazumdar S, Buysse DJ, et al. (1997). Benzodiazepines and zolpidem for chronic insomnia: a meta-analysis of treatment efficacy. *JAMA* 278: 2170–2177.
- Ohayon MM, Roth T (2002). Prevalence of restless legs syndrome and periodic limb movement disorder in the general population. *J Psychosom Res* 53: 547–554.
- Riemann D, Backhaus J (Hrsg.) (1996). Bewältigung von Schlafstörungen. Weinheim: PVU.
- Riemann D, Perlis ML (2009). The treatments of chronic insomnia: a review of benzodiazepine receptor agonists and psychological and behavioral therapies. *Sleep Med Rev* 13: 205–214.
- Riemann D, Voderholzer U (2003). Primary insomnia: a risk factor to develop depression? *J Affect Disord* 76: 255–259.
- Riemann D, Voderholzer U, Cohrs S, et al. (2002). Trimipramine in primary insomnia: results of a polysomnographic double-blind controlled study. *Pharmacopsychiatry* 35: 165–174.

- Riemann D, Spiegelhalder K, Feige B, et al. (2010). The hyperarousal model of insomnia: a review of the concept and its evidence. *Sleep Med Rev* 14: 19–31.
- Schenck CH, Bundlie SR, Ettinger MG, Mahowald MW (1986). Chronic behavioral disorders of human REM sleep: a new category of parasomnia. *Sleep* 9: 293–308.
- Schlack R, Hapke U, Maske U et al. (2013). Häufigkeit und Verteilung von Schlafproblemen und Insomnie in der deutschen Erwachsenenbevölkerung. *Bundesgesundheitsblatt Gesundheitsforschung, Gesundheitsschutz* 5: 740–748.
- Schmidt-Nowara W, Lowe A, Wiegand L, et al. (1995). Oral appliances for the treatment of snoring and obstructive sleep apnea: a review. *Sleep* 18: 501–510.
- Schwartz JR, Roth T (2006). Shift work sleep disorder: burden of illness and approaches to management. *Drugs* 66: 2357–2370.
- Sivertsen B, Omvik S, Pallesen S, et al. (2006). Cognitive behavioral therapy vs. zopiclone for treatment of chronic primary insomnia in older adults: a randomized controlled trial. *JAMA* 295: 2851–2858.
- Smith MT, Perlis ML, Park A, et al. (2002). Comparative meta-analysis of pharmacotherapy and behavior therapy for persistent insomnia. *Am J Psychiatry* 159: 5–11.
- Spoormaker VI, Schredl M, van den Bout J (2006). Nightmares: from anxiety symptom to sleep disorder. *Sleep Med Rev* 10: 19–31.
- Trauer JM, Qian MY, Doyle JS, et al. (2015). Cognitive behavioral therapy for chronic insomnia: a systematic review and meta-analysis. *Ann Intern Med* 163: 191–204.
- Trenkwalder C, Paulus W (2010). Restless legs syndrome: pathophysiology, clinical presentation and management. *Nat Rev Neurol* 6: 337–346.
- Trenkwalder C, Paulus W, Walters AS (2005). The restless legs syndrome. *Lancet Neurol* 4: 465–475.
- Walsh JK, Erman M, Erwin M, et al. (1998). Subjective hypnotic efficacy of trazodone and zolpidem in DMS-III-R primary insomnia. *Hum Psychopharmacol* 13: 191–198.
- Wilt TJ, MacDonald R, Ouellette J, et al. (2013). Pharmacologic therapy for primary restless legs syndrome: a systematic review and meta-analysis. *JAMA Intern Med* 173(7): 496–505.
- Winkler A, Auer C, Doering BK, Rief W (2014). Drug treatment of primary insomnia: a meta-analysis of polysomnographic randomized controlled trials. *CNS Drugs* 28: 799–816.
- Wyatt JK (2004). Delayed sleep phase syndrome: pathophysiology and treatment options. *Sleep* 27: 1195–1203.

19.4 SCHLAFSTÖRUNGEN IM RAHMEN EINER ANDEREN PSYCHIATRISCHEN STÖRUNG

- Benca RM, Obermeyer WH, Thisted RA, Gillin JC (1992). Sleep and psychiatric disorders. A meta-analysis. *Arch Gen Psychiatry* 49: 651–668.
- Baglioni C, Nanovska S, Regen W, et al. (2016). Sleep and mental disorders: a meta-analysis of polysomnographic research. *Psychol Bull* 142: 969–990.

19.5 ANDERE SCHLAFSTÖRUNGEN

- Lader M, Cardinali DP, Pandi-Perumal SR (2006). *Sleep and Sleep Disorders: A Neuropsychopharmacological Approach*. New York: Springer.

LEITLINIEN

- AWMF-S3-Leitlinie Nicht erholsamer Schlaf – Schlafstörungen (2016). AWMF-Registrierungsnummer 063–001. www.awmf.org/leitlinien/detail/II/063-001.html (letzter Zugriff: 20.4.2018).
- American Academy of Sleep Medicine – Guidelines. www.aasmnet.org/practiceguidelines.aspx (letzter Zugriff: 20.4.2018).
- Riemann D, Baum E, Cohrs S et al. (2017). S3-Leitlinie Nicht erholsamer Schlaf/Schlafstörungen. Kapitel „Insomnie bei Erwachsenen“ (AWMF-Registrierungsnummer 063-003), Update 2016. *Somnologie* 21: 2–44.

SYSTEMATISCHE COCHRANE REVIEWS (WWW.COCHRANE.DE/DEUTSCH)

- Giles TL, Lasserson TJ, Smith BH, et al. (2006). Continuous positive airways pressure for obstructive sleep apnea in adults. *Cochrane Database Syst Rev* 3: CD001106.
- Hershner A, Petrie KJ (2002). Melatonin for the prevention and treatment of jet lag. *Cochrane Database Syst Rev* 2: CD001520.
- Mason M, Welsh EJ, Smith I (2013). Drug therapy for obstructive sleep apnoea in adults. *Cochrane Database Syst Rev* 5: CD003002.
- Montgomery P, Dennis J (2002). Bright light therapy for sleep problems in adults aged 60+. *Cochrane Database Syst Rev* 2: CD003403.
- Montgomery P, Dennis J (2003). Cognitive behavioural interventions for sleep problems in adults aged 60+. *Cochrane Database Syst Rev* 1: CD003161.
- Scholz H, Trenkwalder C, Kohnen R, et al. (2011a). Levodopa for restless legs syndrome. *Cochrane Database Syst Rev* 2: CD005504.
- Scholz H, Trenkwalder C, Kohnen R, et al. (2011b). Dopamine agonists in restless legs syndrome. *Cochrane Database Syst Rev* 3: CD006009.
- Shneerson J, Wright J (2001). Lifestyle modification for obstructive sleep apnoea. *Cochrane Database Syst Rev* 1: CD002875.
- Sundaram S, Bridgman SA, Lim J, Lasserson TJ (2005). Surgery for obstructive sleep apnoea. *Cochrane Database Syst Rev* 4: CD001004.
- Trotti LM, Bhadriraju S, Becker LA (2012). Iron for restless legs syndrome. *Cochrane Database Syst Rev* 5: CD007834.
- Vignatelli L, D'Alessandro R, Candelise L (2008). Antidepressant drugs for narcolepsy. *Cochrane Database Syst Rev* 1: CD003724.
- Wozniak DR, Lasserson TJ, Smith I (2014). Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev* 1: CD007736.

PATIENTENRATGEBER

- Patientenratgeber der Deutschen Gesellschaft für Schlafforschung und Schlafmedizin. www.charite.de/dgsm/dgsm/patienteninformationen_ratgeber.php?language=german (letzter Zugriff: 20.4.2018).
- Riemann D (2004). *Ratgeber Schlafstörungen*. Göttingen: Hogrefe.